arXiv:1612.00546v1 [nlin.SI] 2 Dec 2016

An AB-KdV Equation, Exact Solutions, Symmetry Reductions and Bäcklund Transformations

Man JIA[1,2], **Sen Yue LOU**[1,2,3]

[1] Physics Department, Ningbo University, Ningbo 315211, China
[2] Ningbo Collaborative Innovation Center of Nonlinear Hazard System of Ocean and Atmosphere, Ningbo University, Ningbo 315211, China
[3] Shanghai Key Laboratory of Trustworthy Computing, East China Normal University, Shanghai 200062, China

E-mail: jiaman@nbu.edu.cn

Abstract. We study a nonlocal AB-KdV system obtained directly from the KdV equation to describe two-area physical event. The exact solutions of the AB-KdV system, including P_sT_d invariant and P_sT_d symmetric breaking solutions are shown by different methods. These solutions, such as single soliton solutions, infinitely many singular soliton solutions, cnoidal wave solutions, and symmetry reduction solutions etc., show the AB-KdV system possesses rich structures. Also, a special Bäcklund transformation related to residual symmetry is presented via the localization of the residual symmetry to find interactive solutions between the solitons and other types of the AB-KdV system.

1. Introduction

2013, Mark J. Ablowitz [1] introduced a nonlinear Schrödinger equation

$$iq_t(x,t) = q_{xx}(x,t) \pm 2q(x,t)q^*(-x,t)q(x,t), \tag{1}$$

where $*$ denotes complex conjugation and $q(x,\ t)$ is a complex valued function of the real variables x and t. Some of the important properties of the nonlocal NLS equation eq.(1) were contrasted with the classical NLS equation where the nonlocal nonlinear term $q^*(-x,t)$ is replaced by $q^*(x,t)$. They pointed out that both eq. (1) and the classical NLS share the symmetry that when $x \to -x$, $t \to -t$ and a complex conjugate is taken, then the equation remains invariant. Thus, the new nonlocal equation is PT symmetric.

Note that eq. (1) is invariant neither under parity P, whose effect is to make spatial reflections, $p \to -p$ and $x \to -x$, nor under time reversal T , which replaces $p \to -p$, $x \to -x$, and $i \to -i$. And there are other types of nonlocal nonlinear systems, such as the coupled nonlocal NLS systems [2], the nonlocal modified KdV systems [3]-[4], the discrete nonlocal NLS system [5], and the nonlocal Davey-Stewartson system [6]-[8], etc.

PT symmetry is important in not only particle physics and quantum physics [9], but also many other research fields in physics, such as optics [10], quantum field theory [11], electric circuits [12]. Recently, PT-symmetric nonlocal NLS equation (1) is used to describe the extension of properties of traditional macroscopic magnetic systems [13].

In fact, there are other possible crucial properties in particle physics, such as charge conjugation (C), shifted-parity (P_s, parity with a shift, or $x \to -x + x_0$, where x_0 is an arbitrary constant), delayed time reversal (T_d, time reversal with a delay, or $t \to -t+t_0$, where t_0 is an arbitrary constant) and their possible combinations such as PT, PC, P_sC, P_sT_d which can be successively used to describe two-place physical problems. These systems are named Alice-Bob (AB) systems [14] introduced by the AB-BA principle and $P_s - T_d - C$ principle. The nonlocal NLS equation (1) can be considered as a special type of the AB systems.

In [14], some special types of AB systems such as the KdV-KP-Toda type, mKdV-sG type, NLS type and discrete H_1 type are established from the well-known integrable systems. Contrasted to the nonlocal NLS equation (1) solved by inverse scatting method, some special $P_s - T_d - C$ group invariant multi-soliton solutions of the AB systems are obtained by using the $P_s - T_d - C$ principle. Furthermore, in [15] Lou established a general AB-KdV equation and obtained many types of P_sT_d invariant solutions of the general AB-KdV equation, such as the Painlevé II reduction and soliton-cnoidal periodic wave interaction solutions.

Except for the P_sT_d invariant solutions, there exist other types of multiple soliton solutions of P_sT_d symmetry breaking for different AB models. A special P_sT_d symmetry breaking solution for AB-KdV equation is listed in [15].

Hence, two quite relevant questions arise: are there any more possible P_sT_d symmetry breaking solutions for the AB systems? How can we find the solutions? The main purpose of our manuscript is to resolve the issues.

In this manuscript, we construct a special AB-KdV system with the P_sT_d principle and find the exact solutions, including P_sT_d invariant and P_sT_d symmetric breaking solutions of the AB-KdV system. The P_sT_d invariant solutions are obtained by using the $P_s - T_d - C$ principle, while the P_sT_d symmetric breaking solutions are solved by the help of a coupled KdV system. We also find the Bäcklund transformation related to residual symmetry via the localization of the residual symmetry which can be used to find interactive solutions between the solitons and other types of the AB-KdV system.

2. An AB-KdV Equation

In this manuscript, we mainly discuss the following AB-KdV equation

$$\begin{aligned} &A_t + A_{xxx} + 3(A+B)(B_x + 3A_x) = 0, \\ &B = \hat{P}_s\hat{T}_d A = A(-x + x_0, -t + t_0). \end{aligned} \tag{2}$$

The AB-KdV equation is established as follows [15]. Substituting $u = \frac{1}{2}(A + B)$ into the common KdV equation

$$u_t + 24uu_x + u_{xxx} = 0, \tag{3}$$

we have

$$A_t + B_t + A_{xxx} + B_{xxx} + 12(A + B)(A_x + B_x) = 0, \tag{4}$$

which can be split into two equations

$$A_t + A_{xxx} + 3(A + B)(B_x + 3A_x) + G(A,\ B) = 0, \tag{5}$$

$$B_t + B_{xxx} + 3(A + B)(A_x + 3B_x) - G(A,\ B) = 0, \tag{6}$$

with $G(A,\ B)$ may be an arbitrary function of A and B. As B is related to A by $B = \hat{P}_s\hat{T}_dA = A(-x + x_0, -t + t_0)$, we can see $\hat{P}_s\hat{T}_d$ acted on the equation (6) lead to the compatibility condition

$$G(A,\ B) = \hat{P}_s\hat{T}_dG(A,\ B),$$

which means the arbitrary function $G(A,\ B)$ should be P_sT_d invariant. There are numerous functions satisfying P_sT_d invariant. But for simplicity, we take $G(A,\ B)$ is zero and the equations (5)- (6) are reduced to one equation (2).

3. Group invariant solutions of the AB-KdV equation

The AB-KdV system (2) is directly derived from the common KdV equation (3), so all the P_sT_d invariant solutions of the KdV equation (3) are also solutions of the AB-KdV system (2). Various exact solutions of the KdV equation (3) have been studied in literature. Thus we can use those known solutions to select out the P_sT_d invariant solutions for the AB-KdV system (2). Here we list three special significant examples.

3.1. P_sT_d-invariant multi-soliton solutions of the AB-KdV equation

For the KdV equation (3), it is well known multiple soliton solutions possesses the form [16]

$$u = \frac{1}{2}(\ln F)_{xx}, \tag{7}$$

$$F = \sum_{\nu} \exp\left(\sum_{j=1}^{N} \nu_j\xi_j + \sum_{1\le j\le l}^{N} \nu_j\nu_l\theta_{jl}\right), \tag{8}$$

where the summation of ν should be done for all permutations of $\nu_i = 0, 1, i = 1, 2, \cdots, N$ and

$$\xi_j = k_jx - k_j^3t + \xi_{0j}, \qquad \exp(\theta_{jl}) = \left(\frac{k_j - k_l}{k_j + k_l}\right)^2. \tag{9}$$

The solution (7) and (8) is not P_sT_d group invariant. KdV equation (3) is space-time translation invariant which makes every soliton being located at anywhere ξ_{0j}. To discuss the P_sT_d group invariant solutions of the AB-KdV equation (2), we rewrite the solutions of (9) as

$$\xi_j = \eta_j + \eta_{0j} - \frac{1}{2}\sum_{i=1}^{j-1}\theta_{ij} - \frac{1}{2}\sum_{i=j+1}^{N}\theta_{ji}, \tag{10}$$

with η_j being

$$\eta_j = k_j(x - \frac{1}{2}x_0) - k_j^3(t - \frac{1}{2}t_0) + \eta_{0j}, \tag{11}$$

then the N-soliton solutions of the KdV equation (3) is

$$u = \frac{1}{2}\left[\ln \sum_{\nu=-1,1} K_\nu \cosh\left(\frac{1}{2}\sum_{j=1}^{N}\nu_j\eta_j\right)\right]_{xx}, \tag{12}$$

where the summation of $\nu = \{\nu_1,\ \nu_2,\ \cdots,\ \nu_N\}$ should be done for all non-dual permutations of $i = 1,\ 2,\ \cdots,\ N$ and

$$K_\nu = \prod_{i>j}(k_i - \nu_i\nu_j k_j). \tag{13}$$

From the expression (12), it is easy to see that

$$A^{P_sT_d}_{\eta_{0j}=0} = \hat{P}_s\hat{T}_d A_{\eta_{0j}=0} = A_{\eta_{0j}=0}. \tag{14}$$

Then the multiple soliton solutions $A_{\eta_{0j}=0}$ presents the P_sT_d group invariant solutions of the AB-KdV equation (2).

3.2. P_sT_d-invariant symmetry reduction solution of the AB-KdV equation

The KdV equation (3) possesses a special symmetry reduction

$$u = t^{-\frac{2}{3}}U(\xi)^2, \qquad \xi = t^{-\frac{1}{3}}x, \tag{15}$$

where $U = U(\xi)$ satifies

$$3UU_{\xi\xi\xi} - U^2 + 9U_\xi U_{\xi\xi} + 72U^3U_\xi - \xi UU_\xi = 0. \tag{16}$$

In order to satisfy P_sT_d invariant condition for the AB-KdV system (2), we have to rewrite the function U and ξ of the symmetry reduction (15)-(16) of the KdV equation (3), so that the symmetry reduction of the AB-KdV system (2) is

$$\begin{aligned} A &= \left(t - \frac{t_0}{2}\right)^{-\frac{2}{3}} U_1(\xi_1)^2, \\ \xi_1 &= \left(x - \frac{x_0}{2}\right)\left(t - \frac{t_0}{2}\right)^{-\frac{1}{3}}, \end{aligned} \tag{17}$$

where $U_1(\xi_1)$ is a solution of

$$\begin{aligned} &3U_1U_{1\xi\xi\xi} - U_1^2 + 9U_{1\xi_1}U_{1\xi_1\xi_1} + 72U_1^3U_{1\xi_1} \\ &\quad - \xi_1U_1U_{\xi_1} = 0. \end{aligned} \tag{18}$$

3.3. P_sT_d-invariant soliton-cnoidal wave interaction solutions of the AB-KdV equation

The KdV equation (3) has a soliton-cnoidal periodic wave interaction solution [17]

$$u = -\frac{1}{2}w_x^2\tanh(w)^2 + \frac{1}{2}w_{xx}\tanh(w) + \frac{1}{3}w_x^2 - \frac{1}{6}\frac{w_{xxx}}{w_x} - \frac{1}{24}\frac{w_x}{w_t} + \frac{1}{8}\frac{w_{xx}^2}{w_x^2}, \tag{19}$$

$$w = \frac{1}{2}\text{arctanh}(m\text{sn}(k(x-c_2t),m)) + \frac{1}{2}k(x-c_1t), \tag{20}$$

$$k = \sqrt{\frac{c_1-c_2}{2(1-m^2)}}, \tag{21}$$

with arbitrary constants c_1, c_2 and m.

So the P_sT_d invariant soliton-cnoidal periodic wave interaction solution of the AB-KdV system (2) is

$$A = -\frac{1}{2}w_{1x}^2\tanh(w_1)^2 + \frac{1}{2}w_{1xx}\tanh(w_1) + \frac{1}{3}w_{1x}^2 - \frac{1}{6}\frac{w_{1xxx}}{w_{1x}} - \frac{1}{24}\frac{w_{1x}}{w_{1t}} + \frac{1}{8}\frac{w_{1xx}^2}{w_{1x}^2}, \tag{22}$$

$$w_1 = \frac{1}{2}k\xi_1 + \frac{1}{2}\text{arctanh}(m\text{sn}(k\xi_2,m)),$$
$$k = \sqrt{\frac{c_1-c_2}{2(1-m^2)}}, \tag{23}$$

$$\xi_1 = \left(x - \frac{x_0}{2}\right) - c_1\left(t - \frac{t_0}{2}\right),$$
$$\xi_1 = \left(x - \frac{x_0}{2}\right) - c_2\left(t - \frac{t_0}{2}\right), \tag{24}$$

with arbitrary constants c_1, c_2 and m.

4. P_sT_d Symmetry Breaking Solutions of the AB-KdV System

In section III, many P_sT_d-invariant solutions of the AB-KdV system (2), including multiple soliton solutions, P_sT_d-invariant symmetry reduction solutions and soliton-cnoidal wave interaction solutions are found. Then our question is, how to find P_sT_d symmetry breaking solutions of the AB-KdV system? It is also interesting but difficult to find P_sT_d symmetry breaking solutions.

In order to obtain the P_sT_d symmetry breaking soliton solutions of the AB-KdV system (2), we start from a coupled KdV system

$$u_t + u_{xxx} + 3(u+v)(v_x+3u_x) = 0, \tag{25}$$
$$v_t + v_{xxx} + 3(u+v)(u_x+3v_x) = 0, \tag{26}$$

with $u = u(x,t)$ and $v = v(x,t)$ being a normal coupled KdV system. Compared the cKdV system (25)-(26) with the AB-KdV system (2), we know if v is replaced by $u(-x,\ -t)$, the coupled KdV system (25) will be reduced to one equation

$$\begin{aligned} u_t + u_{xxx} + 3[u + u(-x,-t)][u(-x,-t)_x \\ + 3u_x] = 0, \end{aligned} \tag{27}$$

which is a special case of the AB-KdV system (2) with $x_0 = t_0 = 0$. It implies that looking for the exact solutions of the AB-KdV equation (2) is equivalent to find the solutions of the coupled KdV equation (25)-(26). The solutions of the cKdV system (25)-(26) may be changed to those of the AB-KdV (2) by using $v = u(-x,-t)$ due to the P_sT_d symmetric invariant of the AB-KdV equation (2).

In this section, we will show details on how to obtain the P_sT_d symmetric breaking solutions of the AB-KdV system (2), including single soliton solutions, singular soliton solutions and cnoidal wave solutions with the help of the coupled KdV system (25)-(26).

4.1. single soliton solutions of the AB-KdV equation

To find the single soliton solutions of the cKdV system (25)-(26), by means of the tanh function expansion approach, we assume the single soliton solution of the cKdV system (25)-(26) has the form [18]

$$\begin{aligned} u &= a_0 + a_1 \tanh(kx+ct) + a_2 \tanh(kx+ct)^2, \\ v &= b_0 + b_1 \tanh(kx+ct) + b_2 \tanh(kx+ct)^2. \end{aligned} \tag{28}$$

Substituting (28) into the cKdV system (25)-(26) and vanishing all the coefficients of $\tanh(kx+ct)^i$ for i, then determining the arbitrary constant a_0, a_1, b_0, b_1 and c, the single soliton solution of the the cKdV system (25)-(26) reads as

$$\begin{aligned} u &= a_1 \tanh(kx-4k^3t) - \frac{1}{2}k^2 \tanh(kx-4k^3t)^2 \\ &\quad + k^2 - b_0, \\ v &= -a_1 \tanh(kx-4k^3t) - \frac{1}{2}k^2 \tanh(kx-4k^3t)^2 \\ &\quad + b_0, \end{aligned} \tag{29}$$

where a_1 and b_0 are arbitrary constant.

Now we consider the AB-KdV equation (2) where B is linked to A with $B = \hat{P}_s\hat{T}_dA = A(-x+x_0,\ -t+t_0)$. If $x_0 = t_0 = 0$, B is linked to A with $B = A(-x,\ -t)$. Letting $v = u(-x,\ -t)$ in the single soliton solution (29) of the cKdV system (25)- (26), we have

$$\begin{aligned} u(-x,-t) = &- a_1 \tanh(kx-4k^3t) \\ &- \frac{1}{2}k^2 \tanh(kx-4k^3t)^2 + b_0. \end{aligned} \tag{30}$$

It can be easily found that $k^2 - b_0$ should be equal to b_0 which means b_0 is fixed as $b_0 = \frac{k^2}{2}$. So the single soliton solution of AB-KdV equation (2) is

$$\begin{aligned} A &= \frac{k^2}{2} + a_1 \tanh(w) - \frac{1}{2}k^2 \tanh(w)^2, \\ w &= k\left(x - \frac{x_0}{2}\right) - 4k^3\left(t - \frac{t_0}{2}\right), \end{aligned} \tag{31}$$

with a_1 being arbitrary constant.

The single soliton solution (31) is P_sT_d symmetric breaking because of the existence of $\tanh(w)$ term which makes $\hat{P}_s\hat{T}_dA \neq A$, so the single soliton solution (31) of the AB-KdV system (2) is P_sT_d symmetric breaking solution.

4.2. infinitely many singular soliton solutions

Furthermore, we try to discuss the infinitely many soliton solutions of the cKdV equation (25)-(26) by substituting

$$\begin{aligned} u &= a_1U(\eta, t) - \frac{1}{2}k^2\eta^2 + k^2 - b_0, \\ v &= -a_1V(\eta, t) - \frac{1}{2}k^2\eta^2 + b_0, \end{aligned} \tag{32}$$

where $\eta \equiv \tanh(kx - 4k^3t)$ into the cKdV system (25)-(26) to determine the functions $U(\eta, t)$ and $V(\eta, t)$. The determined equations of $U(\eta, t)$ and $V(\eta, t)$ are

$$\begin{aligned} U_t = {} & 3k(\eta^2-1)[3U + 3V + k^2(\eta^2-1)]U_\eta \\ & + 3k(\eta^2-1)[U + V + k^2(\eta^2-1)]V_\eta \\ & + 6\eta k^3(\eta^2-1)^2U_{\eta\eta} + k^3(\eta^2-1)^3U_{\eta\eta\eta} \\ & - 12\eta k^3(\eta^2-1)(U+V), \\ V_t = {} & 3k(\eta^2-1)[3U + 3V + k^2(\eta^2-1)]V_\eta \\ & + 3k(\eta^2-1)[U + V + k^2(\eta^2-1)]U_\eta \\ & + 6\eta k^3(\eta^2-1)^2V_{\eta\eta} + k^3(\eta^2-1)^3V_{\eta\eta\eta} \\ & - 12\eta k^3(\eta^2-1)(U+V). \end{aligned} \tag{33}$$

Once the system (33) is solved, we can obtain the exact solutions of the cKdV equation (25)-(26). Here we don't show it in detail.

To present the infinitely many soliton solutions of the AB-KdV equation (2), we also require the functions satisfy $U(\eta, t) = -V(\eta, t)$, then the coupled system of $U(\eta, t)$ and $V(\eta, t)$ (33) is reduced to a linear one equation

$$V_t = 6k^3\eta(\eta^2-1)^2V_{\eta\eta} + k^3(\eta^2-1)^3V_{\eta\eta\eta}, \tag{34}$$

which can be solved by means of the variable separation approach. A special solution of (34) is

$$V = \sum_{i=1}^{N} a_i \left\{ \left(\frac{1-\eta}{1+\eta}\right)^{c_i} (\eta + c_i) \exp\left[8c_ik^3(c_i^2-1)t\right] \right.$$

$$+\left(\frac{1-\eta}{1+\eta}\right)^{-c_i}(\eta-c_i)\exp[-8c_ik^3(c_i^2-1)t]\Bigg\}$$
$$+c_{01}\eta, \tag{35}$$

with a_i, c_i and c_{01} being arbitrary constant.

As $U(\eta,t)$ and $V(\eta,t)$ are known, using the ansatz $v=u(-x,\ -t)$ in (32), we obtain the infinitely many soliton solution of the AB-KdV equation (2)

$$\begin{aligned}A=&\sum_{i=1}^{N}\left[\left(\frac{1-\eta_1}{1+\eta_1}\right)^{c_i}(\eta_1+c_i)\exp[8c_ik^3(c_i^2-1)t]\right.\\&\left.+\left(\frac{1-\eta_1}{1+\eta_1}\right)^{-c_i}(\eta_1-c_i)\exp[-8c_ik^3(c_i^2-1)t]\right]\\&+c_{01}\eta_1-\frac{1}{2}k^2\eta_1^2+\frac{1}{2}k^2,\\\eta_1=&\tanh\left[k\left(x-\frac{x_0}{2}\right)-4k^3\left(t-\frac{t_0}{2}\right)\right].\end{aligned}\tag{36}$$

It is interesting that the solution (36) is not only singularity, but also P_sT_d symmetric breaking because the existence of $c_{01}\eta_1$.

4.3. cnoidal wave solutions

Except for the soliton solutions, we can also try to explore the cnoidal wave solutions of the cKdV system (25)-(26) and AB-KdV equation (2). Like the procedure of searching for soliton solutions, we first consider the cnoidal wave solutions of the cKdV system (25)-(26), then try to find the connections between the cKdV system and AB-KdV equation, and apply the relationship to present the solutions of the AB-KdV equation.

Using the elliptic function expansion method, it is natural to assume the cKdV system (25)-(26) possesses the cnoidal wave solution in Jacobi elliptic function form [19]

$$\begin{aligned}u&=a_0+a_1\mathrm{sn}(kx+ct,m)+a_2\mathrm{sn}(kx+ct,m)^2,\\v&=b_0+b_1\mathrm{sn}(kx+ct,m)+b_2\mathrm{sn}(kx+ct,m)^2,\end{aligned}\tag{37}$$

where a_0, a_1 b_0, b_1 and c are constant to be determined. Substituting (37) into the cKdV system (25)-(26) and vanishing all the coefficients of $\mathrm{sn}(kx+ct,m)^i$ for different i, the exact solutions of (25)-(26) can be read as

$$\begin{aligned}u=&a_1\mathrm{sn}[kx-2k^3(m^2+1)t,m]\\&-\frac{1}{2}k^2\mathrm{sn}[kx-2k^3(m^2+1)t,m]^2+a_0,\\v=&-a_1\mathrm{sn}[kx-2k^3(m^2+1)t,m]\\&-\frac{1}{2}k^2\mathrm{sn}[kx-2k^3(m^2+1)t,m]^2\\&+\frac{1}{2}k^2m^2+\frac{1}{2}k^2-a_0,\end{aligned}\tag{38}$$

with a_0 and a_1 being arbitrary constant.

To obtain the exact solutions of the AB-KdV equation (2), we can let $v = u(-x, -t)$ in the cKdV system (25)-(26) which makes cKdV system (26) reduce to a sepcial AB-KdV equation (2) with B linked with A in the form of $B = A(-x, -t)$. Letting $v = u(-x, -t)$ in (38) leads to

$$a_0 = \frac{1}{2}k^2m^2 + \frac{1}{2}k^2 - a_0, \tag{39}$$

which means a_0 should be determined as

$$a_0 = \frac{1}{4}k^2m^2 + \frac{1}{4}k^2. \tag{40}$$

Then the corresponding cnoidal wave solution of AB-KdV equation (2) is

$$\begin{aligned} A &= a_1\zeta - \frac{1}{2}k^2\zeta^2 + \frac{1}{4}k^2m^2 + \frac{1}{4}k^2, \\ \zeta &= \mathrm{sn}\left[k\left(x - \frac{x_0}{2}\right) - 2k^3(m^2+1)\left(t - \frac{t_0}{2}\right), m\right], \end{aligned} \tag{41}$$

with a_1, k and m being arbitrary constant. The cnoidal wave solution (41) of the AB-KdV system (2) in Jacobi elliptic form is also anti-P_sT_d symmetric.

4.4. Infinitely many periodic wave solutions

In order to obtain the infinitely many periodic wave solutions of the AB-KdV equation, we try to find the infinitely many periodic solutions of the cKdV equation (25)-(26) first. Substituting

$$\begin{aligned} u &= F(\omega, t) - \frac{1}{2}k^2m^2\omega^2 + a_0, \\ v &= G(\omega, t) - \frac{1}{2}k^2m^2\omega^2 + \frac{1}{2}k^2m^2 + \frac{1}{2}k^2 - a_0, \end{aligned} \tag{42}$$

where $\omega \equiv \mathrm{sn}[kx - 2k^3(m^2+1)t, m]$, $F(\omega, t)$ and $G(\omega, t)$ are the functions of (ω, t) into the cKdV equation (25)-(26), we can find the determining equations of $F(\omega, t)$ and $G(\omega, t)$ read as

$$\begin{aligned} F_t = &-\frac{3}{2}z_0k(6F + 6G + k^2z_1)F_\omega - \frac{3}{2}z_0k(2F + 2G \\ &+ k^2z_1)G_\omega - 3z_0z_1k^3\omega F_{\omega\omega} - k^3z_0^3F_{\omega\omega\omega} \\ &+ 12z_0\omega m^2k^3(F + G), \\ G_t = &-\frac{3}{2}z_0k(6F + 6G + k^2z_1)G_\omega - \frac{3}{2}z_0k(2F + 2G \\ &+ k^2z_1)F_\omega - 3z_0z_1k^3\omega G_{\omega\omega} - k^3z_0^3G_{\omega\omega\omega} \\ &+ 12z_0\omega m^2k^3(F + G), \end{aligned} \tag{43}$$

with

$$\begin{aligned} z_0 &= \sqrt{1-\omega^2}\sqrt{1-m^2\omega^2}, \\ z_1 &= m^2 - 2m^2\omega^2 + 1. \end{aligned} \tag{44}$$

To search for the solutions of $F(\omega, t)$ and $G(\omega, t)$ of (43) with (44), it is natural to assume the simplest relation of F and G satisfying $F(\omega, t) = -G(\omega, t) + H(\omega, t)$ for any

given solutions of $H(\omega,t)$. Under the condition of $F(\omega,t)=-G(\omega,t)+H(\omega,t)$, the determining equations (43) become

$$\begin{aligned} H_t=&-3z_0k(4H+k^2z_1)H_\omega-3z_0z_1k^3\omega H_{\omega\omega}-k^3z_0^3H_{\omega\omega\omega}\\ &+24z_0\omega m^2k^3H, \end{aligned} \tag{45}$$

and especially for the G part, the equation is

$$\begin{aligned} G_t=&-\frac{3}{2}z_0k(2H+z_1)H_\omega-6kz_0HG_\omega-3z_0z_1\omega k^3G_{\omega\omega}\\ &-k^3z_0^3G_{\omega\omega\omega}+12z_0\omega m^2k^3H, \end{aligned} \tag{46}$$

which is a linear one for G with z_0 and z_1 satisfying (44). Eqs. (45) and (46) are still difficult to solve, but we know the equation (45) possesses a special solution of $H(\omega,t)=0$, and then the G part equation is

$$G_t=-3z_0z_1\omega k^3G_{\omega\omega}-k^3z_0^3G_{\omega\omega\omega}, \tag{47}$$

which can be solved by means of the variable separation approach. A special solution of (47) is

$$\begin{aligned} G=&\frac{C_1\omega}{m^2-1}\mathrm{EllipticF}(\omega,m)+\frac{C_1(m^2+1)\omega}{(m^2-1)^2}\mathrm{EllipticE}(\omega,m)\\ &+\frac{C_1(m^2+1)}{(m^2-1)^2}\sqrt{(1-\omega^2)(1-m^2\omega^2)}+C_2\omega+C_3, \end{aligned} \tag{48}$$

where $\mathrm{EllipticF}(\omega,m)$ and $\mathrm{EllipticE}(\omega,m)$ are incomplete elliptic integrals of the first kind and second kind respectively, defined as

$$\mathrm{EllipticF}(\omega,m)=\int_0^\omega\frac{1}{\sqrt{1-t^2}\sqrt{1-m^2t^2}}\mathrm{d}t,$$

$$\mathrm{EllipticE}(\omega,m)=\int_0^\omega\frac{\sqrt{1-m^2t^2}}{\sqrt{1-t^2}}\mathrm{d}t,$$

and C_1, C_2 and C_3 are arbitrary constance. When G is solved, we can obtain F by $F=-G$, then the infinitely many periodic wave solutions of the cKdV equation (25)-(26) are obtained.

As the exact solutions of the cKdV equation (25)-(26) are obtained, we can rewrite the solutions to satisfy the AB-KdV system by $v=u(-x,-t)$. Here we write down the results.

The AB-KdV system possesses the infinitely many periodic wave solutions as follow

$$\begin{aligned} u=&-G_1(\omega_1,t)+H_1(\omega_1,t)-\frac{1}{2}k^2m^2\omega_1^2+\frac{1}{4}k^2m^2+\frac{1}{4}k^2,\\ v=&G_1(\omega_1,t)-\frac{1}{2}k^2m^2\omega_1+\frac{1}{4}k^2m^2+\frac{1}{4}k^2,\\ \omega_1=&\mathrm{sn}\left[k\left(x-\frac{x_0}{2}\right)-2k^3(m^2+1)\left(t-\frac{t_0}{2}\right),m\right], \end{aligned} \tag{49}$$

with $G_1(\omega_1,t)$ and $H_1(\omega_1,t)$ satisfying

$$\begin{aligned} H_{1t}=&-3z_0k(4H_1+k^2z_1)H_{1\omega_1}-3z_0z_1k^3\omega H_{1\omega_1\omega_1}\\ &-k^3z_0^3H_{1\omega_1\omega_1\omega_1}+24z_0\omega_1m^2k^3H_1, \end{aligned} \tag{50}$$

$$
\begin{aligned}
G_{1t} = & -\frac{3}{2}z_0k(2H_1+z_1)H_{1\omega_1} - 6kz_0H_1G_{1\omega_1} \\
& - 3z_0z_1\omega_1k^3G_{\omega_1\omega_1} - k^3z_0^3G_{1\omega_1\omega_1\omega_1} \\
& + 12z_0\omega_1m^2k^3H_1. \qquad (51)
\end{aligned}
$$

We believe that there are possible more soliton solutions and periodic wave solutions with (50) and (51).

A special solution of the AB-KdV system is

$$
\begin{aligned}
u &= -G_1(\omega_1,t) - \frac{1}{2}k^2m^2\omega_1^2 + \frac{1}{4}k^2m^2 + \frac{1}{4}k^2, \\
v &= G_1(\omega_1,t) - \frac{1}{2}k^2m^2\omega_1 + \frac{1}{4}k^2m^2 + \frac{1}{4}k^2, \\
\omega_1 &= \mathrm{sn}\left[k\left(x-\frac{x_0}{2}\right) - 2k^3(m^2+1)\left(t-\frac{t_0}{2}\right), m\right], \qquad (52)
\end{aligned}
$$

and G_1 is the solution of

$$
G_{1t} = -3z_0z_1\omega_1k^3G_{1\omega_1\omega_1} - k^3z_0^3G_{1\omega_1\omega_1\omega_1}, \qquad (53)
$$

and

$$
\begin{aligned}
z_0 &= \sqrt{1-\omega_1^2}\sqrt{1-m^2\omega_1^2}, \\
z_1 &= m^2 - 2m^2\omega^2 + 1. \qquad (54)
\end{aligned}
$$

We should mention that all the anti-symmetric solutions of (53) and (54) lead to the P_sT_d symmetry breaking solutions of the AB-KdV system (2).

5. Lie Point Symmetry and Symmetry Reduction

The knowledge of the symmetries is very useful to enhance our understanding of complex physical phenomena, to simplify and even completely solve the complicated problems. Furthermore, the study of symmetries has been manifested as one of the most important and powerful methods in almost every branch of science especially in physics and mathematics. It is particularly fundamental to find the symmetries of a nonlinear equation in the development of the theory of the integrable systems because of the existence of infinitely many symmetries [20]- [22]. Recent studies show that symmetry reduction approach with nonlocal symmetries and residual symmetries has been successfully used to find some types of interaction solutions of nonlinear excitations for a number of integrable system [23]- [25]. In this section, we will discuss the Lie point symmetry and symmetry reduction solutions of the AB-KdV equation (2) with the help of the cKdV system (25)-(26).

A symmetry, σ is defined by a solution of its linearized equation. For the cKdV system (25)-(26), its symmetries are solutions of

$$
\begin{aligned}
& \sigma_t^u + \sigma_{xxx}^u + 3(\sigma^u+\sigma^v)(v_x+3u_x) + 3(u \\
& \quad + v)(\sigma_x^v + 3\sigma_x^u) = 0, \\
& \sigma_t^v + \sigma_{xxx}^v + 3(\sigma^u+\sigma^v)(u_x+3v_x) + 3(u \\
& \quad + v)(\sigma_x^u + 3\sigma_x^v) = 0. \qquad (55)
\end{aligned}
$$

That means the cKdV system (25)-(26) is form-invariant under the transformation

$$\begin{pmatrix} u \\ v \end{pmatrix} \to \begin{pmatrix} u+\epsilon\sigma^u \\ v+\epsilon\sigma^v \end{pmatrix}, \tag{56}$$

where ϵ is an infinitesimal parameter.

The generalized Lie point symmetries of the cKdV system (25)-(26) possess the form

$$\begin{aligned} \begin{pmatrix} \sigma^u \\ \sigma^v \end{pmatrix} = X(x,t,u,v) \begin{pmatrix} u_x \\ v_x \end{pmatrix} + T(x,t,u,v) \begin{pmatrix} u_t \\ v_t \end{pmatrix} \\ - \begin{pmatrix} U(x,t,u,v) \\ V(x,t,u,v) \end{pmatrix}, \end{aligned} \tag{57}$$

where $X(x,t,u,v)$, $T(x,t,u,v)$, $U(x,t,u,v)$ and $V(x,t,u,v)$ are the functions of the indicated variables.

Substituting (90) into the symmetry definition (55) and using the cKdV system (25)-(26) to eliminate nonindependent quantities, u_t and v_t, we can get a set of determining equations for X, T, U and V. Solving the determining equation system, the Lie point symmetry can be obtained. Here we list the results. The cKdV system (25)-(26) possesses the Lie point symmetries

$$\begin{aligned} \sigma^u &= (c_1x+c_2)u_x + (3c_1t+c_3)u_t + (2c_1+c_4)u \\ &\quad - c_4v - c_0, \\ \sigma^v &= (c_1x+c_2)v_x + (3c_1t+c_3)u_t + (2c_1+c_4)v \\ &\quad - c_4u + c_0, \end{aligned} \tag{58}$$

where c_i, $i=0,1,2,3,4$ are arbitrary constants.

To find symmetry reductions of a nonlinear system means to find the group invariant solutions which is guaranteed by $\sigma^u = \sigma^v = 0$,

$$\begin{aligned} 0 &= (c_1x+c_2)u_x + (3c_1t+c_3)u_t + (2c_1+c_4)u \\ &\quad - c_4v - c_0, \\ 0 &= (c_1x+c_2)v_x + (3c_1t+c_3)u_t + (2c_1+c_4)v \\ &\quad - c_4u + c_0. \end{aligned} \tag{59}$$

Solving the group invariant conditions (59) and the cKdV system (25)-(26), we can get the symmetry reduction solutions. Here we just list a special reduction solution because we aim to find the symmetry reductions of the AB-KdV system (2).

If $\{U(\xi),\ V(\xi)\}$ is the solution of the reduction equation

$$\begin{aligned} U_{\xi\xi\xi} &= \left(\frac{1}{3}\xi - 6V\right)U_\xi - dU, \\ V_{\xi\xi\xi} &= \left(\frac{1}{3}\xi - 12V\right)V_\xi + \frac{2}{3}V, \end{aligned} \tag{60}$$

then

$$\begin{aligned} u &= t^dU + \frac{1}{2}t^{-\frac{2}{3}}V + C_0, \\ v &= -t^dU + \frac{1}{2}t^{-\frac{2}{3}}V - C_0, \end{aligned} \tag{61}$$

is a solution of the cKdV system (25)-(26), where d and C_0 are arbitrary constants with the independent variable $\xi = xt^{-\frac{1}{3}}$.

Now we have obtained the symmetry reduction (60) and (61) of the cKdV system (25)-(26), it is natural for us to utilize the results of cKdV system (25)-(26) to the AB-KdV equation (2) by assuming $v = u(-x, -t)$. And we should keep the P_sT_d invariant of the symmetry reduction to satisfy the AB-KdV equation (2). So the symmetry reduction solution of the AB-KdV equation (2) can be stated in the following theorem.

Theorem 1 (Symmetry reduction theorem of the AB-KdV system (2)). If $\{U_1(\xi_1),\ V_1(\xi_1)\}$ is a solution of the reduction equation

$$\begin{aligned} U_{1\xi_1\xi_1\xi_1} &= \left(\frac{1}{3}\xi_1 - 6V_1\right) U_{1\xi_1} - \frac{2n+1}{2m+1}U_1, \\ V_{1\xi_1\xi_1\xi_1} &= \left(\frac{1}{3}\xi_1 - 12V_1\right) V_{1\xi_1} + \frac{2}{3}V_1, \end{aligned} \tag{62}$$

then

$$\begin{aligned} A &= \left(t - \frac{t_0}{2}\right)^{\frac{2n+1}{2m+1}} U_1 + \frac{1}{2}\left(t - \frac{t_0}{2}\right)^{-\frac{2}{3}} V_1, \\ \xi_1 &= \left(x - \frac{x_0}{2}\right)\left(t - \frac{t_0}{2}\right)^{-\frac{1}{3}}, \end{aligned} \tag{63}$$

with m and n being arbitrary constants, is a solution of the AB-KdV equation (2).

We should point out that the reduction (62) and (63) solution has rich and various structures due to the arbitrary constants m and n. For example, the reduction equation (62) has a zero solution of $V_1 = 0$, with the function U_1 satisfying

$$U_{1\xi_1\xi_1\xi_1} = \frac{1}{3}\xi_1 U_{1\xi_1} - \frac{2n+1}{2m+1}U_1, \tag{64}$$

which has many types of solutions due to the different choice of m and n. For simplicity, we list two examples.

Example 1. If we take $m = n = 0$, then

$$U_1 = C_1(\xi_1^3 - 6), \quad \xi_1 = \left(x - \frac{x_0}{2}\right)\left(t - \frac{t_0}{2}\right)^{-\frac{1}{3}}, \tag{65}$$

with C_1 being arbitrary constant is the polynomial of ξ_1. Substituting (65) and $V_1 = 0$ into (63), we can find the AB-KdV equation (2) possesses a P_sT_d symmetry breaking polynomial solution

$$A = \left(x - \frac{x_0}{2}\right)^3 - 6C_1\left(t - \frac{t_0}{2}\right). \tag{66}$$

Example 2. If we take $m = 0$, $n = -1$, then

$$\begin{aligned} U_1 &= C_2\xi_1^{\frac{3}{2}}\Gamma\left(\frac{2}{3}\right) J_{-\frac{1}{3}}\left(\frac{2\sqrt{3}}{9}\xi_1^{\frac{3}{2}}\right), \\ \xi_1 &= \left(x - \frac{x_0}{2}\right)\left(t - \frac{t_0}{2}\right)^{-\frac{1}{3}}, \end{aligned} \tag{67}$$

where $\Gamma\left(\frac{2}{3}\right)$ is the indicated Γ function and $J_{-\frac{1}{3}}\left(\frac{2\sqrt{3}}{9}\xi_1^{\frac{3}{2}}\right)$ is the Bessel function, and C_2 is arbitrary constant. Substituting (67) and $V_1 = 0$ into (63), we can find the AB-KdV equation (2) has a P_sT_d Bessel solution. Here we don't show it in detail.

6. Bäcklund transformation of the AB-KdV system related to residual symmetry

Residual symmetries have been successfully used to find some types of interaction solutions of nonlinear excitations. Further more, according to the results of the symmetry reductions with nonlocal symmetries, nonlocal residual symmetries can be used to find the Darboux transformations and Bäcklund transformations for some nonlinear systems [23]. In this section, we discuss the residual nonlocal symmetries of the cKdV system (25)-(26), then localize the obtained nonlocal symmetries by introducing an enlarged system to provide the nonlocal symmetries related Bäcklund transformation. At last, we will try to apply the known results to the AB-KdV equation (2).

6.1. Bäcklund transformation theorem related to residual symmetry of the cKdV system

The key point to find residual symmetries is to find the truncated Painlevé expansion of the system. For the cKdV equation (25)-(26), its truncated Painlevé expansion has the form

$$u = \frac{u_0}{f^2} + \frac{u_1}{f} + u_2, \qquad v = \frac{v_0}{f^2} + \frac{v_1}{f} + v_2, \tag{68}$$

where the function f is the singularity manifold, and the functions u_0, u_1, u_2, v_0, v_1 and v_2 are related to the derivatives of f and should be determined by substituting (68) into (25)-(26) and vanishing the coefficients of f^i for different i.

Substituting (68) into the cKdV equation (25)-(26), and vanishing the coefficients of f^i for different i, we can obtain u_0, u_1, u_2, v_0, v_1 and v_2

$$u_0 = -\frac{1}{2}f_x^2, \tag{69}$$

$$u_1 = \frac{1}{2}f_{xx} + q, \tag{70}$$

$$u_2 = -\frac{1}{8}\frac{f_{xxx}}{f_x} + \frac{1}{16}\frac{f_{xx}^2}{f_x^2} + r - \frac{1}{24}\lambda, \tag{71}$$

$$v_0 = -\frac{1}{2}f_x^2, \tag{72}$$

$$v_1 = \frac{1}{2}f_{xx} - q, \tag{73}$$

$$v_2 = -\frac{1}{8}\frac{f_{xxx}}{f_x} + \frac{1}{16}\frac{f_{xx}^2}{f_x^2} - r - \frac{1}{24}\lambda, \tag{74}$$

where q and r satisfy the following equations

$$q_{xx} = -f_x r_x + \frac{f_{xx}q_x}{f_x} - \frac{1}{2}\frac{f_t}{f_x}q + \frac{1}{3}\lambda q, \tag{75}$$

$$q_t = 2f_x r_{xx} - 2f_{xx}r_x + \frac{2}{3}\lambda q_x - \frac{\lambda q f_{xx}}{3} + \frac{f_{xt}q}{f_x}, \tag{76}$$

$$r_t = -r_{xxx} + 2\lambda r_x - \frac{3}{2}\frac{f_t r_x}{f_x} + \frac{3}{2}\frac{f_{xx}^2 r_x}{f_x^2}, \tag{77}$$

with λ being an arbitrary constant and f being a solution of the Shwarzian KdV equation [17]

$$f_t = -f_{xxx} + \frac{3}{2}\frac{f_{xx}^2}{f_x} + \lambda f_x. \tag{78}$$

Now the truncated Painlevé expansion has been given. It can be proved that $\{u_1,\ v_1\}$ in the truncated Painlevé expansion (68) is a symmetry with respect to the solution $\{u_2,\ v_2\}$ of the cKdV system (25)-(26) which is called residue symmetry [23] related to Darboux transformations (DTs) and Bäcklund transformations (BTs). Furthermore, the function f, f_x and f_{xx} is linked with $\{u,\ v\}$ non-locally via (71) and (74). The symmetry $\{u_1,\ v_1\}$ is nonlocal. We can verify it it true by substituting $\sigma^u = u_1$ and $\sigma^v = v_1$ with the solution $\{u_2,\ v_2\}$ of the cKdV system (25)-(26) into the symmetry definition equation (55), where q, r and f satisfy the equations (75)-(78), with λ being an arbitrary constant.

Since $\{u_1,\ v_1\}$ is a non-local symmetry related to DTs and BTs, one can naturally believe that it can be localized to a Lie point symmetry such that we can use the initial value problem to link two solutions of the cKdV system (25)-(26) due to the Lie's first principle. To localization the nonlocal symmetry $\{u_1,\ v_1\}$, we enlarge the cKdV system (25)-(26) into the following system

$$u_t + u_{xxx} + 3(u+v)(v_x + 3u_x) = 0, \tag{79}$$

$$v_t + v_{xxx} + 3(u+v)(u_x + 3v_x) = 0, \tag{80}$$

$$f_t = -f_{xxx} + \frac{3}{2}\frac{f_{xx}^2}{f_x} + \lambda f_x, \tag{81}$$

$$f_1 = f_x, \tag{82}$$

$$f_2 = f_{1x}, \tag{83}$$

$$q_t = 2f_x r_{xx} - 2f_{xx} r_x + \frac{2}{3}\lambda q_x - \frac{\lambda q f_{xx}}{3} + \frac{f_{xt} q}{f_x}, \tag{84}$$

$$q_{xx} = -f_x r_x + \frac{f_{xx} q_x}{f_x} - \frac{1}{2}\frac{f_t}{f_x} q + \frac{1}{3}\lambda q, \tag{85}$$

$$r_t = -r_{xxx} + 2\lambda r_x - \frac{3}{2}\frac{f_t r_x}{f_x} + \frac{3}{2}\frac{f_{xx}^2 r_x}{f_x^2}, \tag{86}$$

$$u = -\frac{1}{8}\frac{f_{xxx}}{f_x} + \frac{1}{16}\frac{f_{xx}^2}{f_x^2} + r - \frac{1}{24}\lambda, \tag{87}$$

$$v = -\frac{1}{8}\frac{f_{xxx}}{f_x} + \frac{1}{16}\frac{f_{xx}^2}{f_x^2} - r - \frac{1}{24}\lambda, \tag{88}$$

$$s_{xx} = \frac{f_{xx}s_x}{f_x} + \left(\frac{1}{2}\frac{f_{xxx}}{f_x} - \frac{3}{4}\frac{f_{xx}^2}{f_x^2} - \frac{1}{6}\lambda\right)s + 4f_x(fr_x - q_x). \tag{89}$$

That means we have to solve the symmetry definition equations

$$\sigma_t^u + \sigma_{xxx}^u + 3(\sigma^u + \sigma^v)(v_x + 3u_x) + 3(u + v)(\sigma_x^v + 3\sigma_x^u) = 0, \tag{90}$$

$$\sigma_t^v + \sigma_{xxx}^v + 3(\sigma^u + \sigma^v)(u_x + 3v_x) + 3(u + v)(\sigma_x^u + 3\sigma_x^v) = 0, \tag{91}$$

$$\sigma_x^f(f_xf_{xxx} + f_xf_t - 3f_{xx}^2) - f_x(f_x\sigma_{xxx}^f + f_x\sigma_t^f - 3f_{xx}\sigma_{xx}^f) = 0, \tag{92}$$

$$\sigma^{f_1} - \sigma_x^f = 0, \tag{93}$$

$$\sigma^{f_2} - \sigma_{1x}^f = 0, \tag{94}$$

$$3f_{xt}(f_x\sigma^q - \sigma_x^f q) + 2f_{xx}(\lambda q\sigma_x^f - \lambda\sigma^q f_2 - 6f_x^2\sigma_x^r) + 12f_x^2r_{xx}\sigma_x^f + 3qf_x\sigma_{xt}^f - 2\sigma_{xx}^f(\lambda qf_x + 6r_xf_x^2) + 12f_x^3\sigma_{xx}^r + 2f_x^2(2\lambda\sigma_x^q - 3\sigma_t^q) = 0, \tag{95}$$

$$3f_tq\sigma_x^f - 12f_x^3\sigma_x^r - 2f_x^2(\lambda q + 6\sigma_x^fr_x + 3\sigma_{xx}^q) - 3f_x(\sigma^qf_t + q\sigma^f - 2f_{xx}\sigma_x^q - 2q_x\sigma_{xx}^f) - 6f_{xx}q_x\sigma_x^f = 0, \tag{96}$$

$$f_x^3(4\lambda\sigma_x^r - 2\sigma_t^r - 2\sigma_{xxx}^r) - 3f_x^2(\sigma_x^rf_t + \sigma_t^rr_x) + 3f_3f_{xx}^2\sigma + 3f_xr_t(f_t\sigma_x^f + 2f_{xx}\sigma_{xx}^f) - 6f_{xx}^2r_x\sigma_x^f = 0, \tag{97}$$

$$f_{xx}^2\sigma_x^f + \sigma_{xxx}^ff_x^2 - f_x(\sigma_{xx}^ff_{xx} + \sigma_x^ff_{xxx}) + 8f_x^3(\sigma^u - \sigma^r) = 0, \tag{98}$$

$$f_{xx}^2\sigma_x^f + \sigma_{xxx}^ff_x^2 - f_x(\sigma_{xx}^ff_{xx} + \sigma_x^ff_{xxx}) + 8f_x^3(\sigma^v + \sigma^r) = 0, \tag{99}$$

$$6\sigma^s(f_xf_{xxx} - 3f_{xx}^2) - 12f_x\sigma^f(4f_x^2fr_x - 4f_x^2q_x - s_xf_{xx}) - 6s(f_x\sigma_{xxx}^f - 3f_{xx}\sigma_{xx}^f) + \sigma^sf_x(2\lambda f_x^2 - 6f_xf_{xxx} + 9f_{xx}^2) - 12f_x^2(4ff_x^2\sigma_x^r + 4f_x^2r_x\sigma^f - 4f_x^2\sigma_x^q - \sigma_{xx}^sf_x + \sigma_x^sf_{xx} + s_x\sigma_{xx}^f) = 0, \tag{100}$$

which are the linearized equations of the system (79)-(89). It is not difficult to verify that the corresponding localized Lie point symmetry of (79)-(89) has the form

$$\sigma^u = \frac{1}{2}f_2 + q, \qquad \sigma^v = \frac{1}{2}f_2 - q, \qquad \sigma^f = -f^2,$$
$$\sigma^{f_1} = -2ff_1, \qquad \sigma^{f_2} = -2ff_2 - 2f_1^2, \qquad \sigma^q = s,$$
$$\sigma^s = -3sf, \qquad \sigma^r = q. \tag{101}$$

The result tells us that the nonlocal symmetry of the cKdV system (25)-(26) becomes a Lie point symmetry for the enlarged system (79)-(89).

Whence a nonlocal symmetry is localized, it can be used to find its finite transformations and the related symmetry reductions. Here we just write down the Bäcklund transformation theorem of the cKdV system (25)-(26) with finite transformations related to the initial value problem. And the corresponding symmetry reductions related to the nonlocal symmetry (101) will be discussed in another manuscript.

Theorem 2 (Bäcklund transformation theorem of the cKdV system (25)-(26)). If $\{u,\ v,\ f,\ f_1,\ f_2,\ q,\ r,\ s\}$ is a solution of the enlarged system (79)-(89), so $\{u',\ v',\ f',\ f_1',\ f_2',\ q',\ r',\ s'\}$ is a group invariant solution with

$$\begin{aligned}
u' &= -\frac{\epsilon^2 f_1^2}{2(1+\epsilon f)^2} + \frac{\epsilon f_2}{2(1+\epsilon f)} + \frac{\epsilon^2 s}{2(1+\epsilon f)} + \epsilon q + u,\\
v' &= -\frac{\epsilon^2 f_1^2}{2(1+\epsilon f)^2} + \frac{\epsilon f_2}{2(1+\epsilon f)} - \frac{\epsilon^2 s}{2(1+\epsilon f)} - \epsilon q + v,\\
f' &= \frac{f}{1+\epsilon f}\\
f_1' &= \frac{f_1}{(1+\epsilon f)^2},\\
f_2' &= \frac{f_2}{(1+\epsilon f)^2} - \frac{2\epsilon f_1^2}{(1+\epsilon f)^3},\\
q' &= \frac{\epsilon(2+\epsilon f)s}{2(1+\epsilon f)^2} + q,\\
s' &= \frac{s}{(1+\epsilon f)^3},\\
r' &= \frac{\epsilon^2 s}{2(1+\epsilon f)} + \epsilon q + r.
\end{aligned} \tag{102}$$

The theorem 2 can be proved by solving the following initial value problem,

$$\begin{aligned}
\frac{\mathrm{d}u'(\epsilon)}{\mathrm{d}\epsilon} &= \frac{1}{2}f_2'(\epsilon) + q'(\epsilon), \qquad u'(0) = u,\\
\frac{\mathrm{d}v'(\epsilon)}{\mathrm{d}\epsilon} &= \frac{1}{2}f_2'(\epsilon) - q'(\epsilon), \qquad v'(0) = v,\\
\frac{\mathrm{d}f'(\epsilon)}{\mathrm{d}\epsilon} &= -f'(\epsilon)^2, \qquad f'(0) = f,\\
\frac{\mathrm{d}f_1'(\epsilon)}{\mathrm{d}\epsilon} &= -2f'(\epsilon)f_1'(\epsilon), \qquad f_1'(0) = f_1,\\
\frac{\mathrm{d}f_2'(\epsilon)}{\mathrm{d}\epsilon} &= -2f'(\epsilon)f_2'(\epsilon) - 2f_1'(\epsilon)^2, \qquad f_2'(0) = f_2,\\
\frac{\mathrm{d}q'(\epsilon)}{\mathrm{d}\epsilon} &= s'(\epsilon), \qquad q'(0) = s,\\
\frac{\mathrm{d}s'(\epsilon)}{\mathrm{d}\epsilon} &= -3s'(\epsilon)f'(\epsilon), \qquad s'(0) = s,
\end{aligned}$$

$$\frac{\mathrm{d}r'(\epsilon)}{\mathrm{d}\epsilon} = q'(\epsilon), \qquad r'(0) = q. \tag{103}$$

According to theorem 2, starting from any seed solutions of the cKdV system (25)-(26), we can obtain infinitely many new solutions, especially interaction solutions among different nonlinear excitations [25]. We will discuss it in another manuscript.

6.2. Bäcklund transformation theorem related to residual symmetry of the AB-KdV system

Based on the known results of the cKdV system (25)-(26), we now consider the Bäcklund transformation for the AB-KdV system (2).

By using $v = u(-x,-t)$ in the cKdV equation (26), the solutions of the cKdV system (25)-(26) may be changed to those of the AB-KdV (2) due to the P_sT_d symmetric invariant of the AB-KdV equation (2). It is easily to verify the following Bäcklund transformation theorem of the AB-KdV system (2) from the cKdV system.

Theorem 3 (Bäcklund transformation theorem for AB-KdV system (2)). If A is solution of the AB-KdV system (2), f is a group invariant solution of the Shwarzian KdV equation,

$$f_t = -f_{xxx} + \frac{3}{2}\frac{f_{xx}^2}{f_x} + \lambda f_x, \tag{104}$$

and $\{s,\ q,\ r\}$ are anti-symmetric ($s(-x,-t) = -s(x,t)$, $q(-x,-t) = -q(x,t)$, $r(-x,-t) = -r(x,t)$) solutions of

$$\begin{aligned} q_t = {} & \frac{1}{2}\frac{f_{tx}}{f_x} - \left(2r_x + \frac{1}{3}\lambda\frac{q}{f_x}\right) f_{xx} + 2r_{xx}f_x \\ & + \frac{2}{3}\lambda q_x, \end{aligned} \tag{105}$$

$$\begin{aligned} q_{xx} = {} & \frac{1}{2}q\frac{f_{xxx}}{f_x} - \frac{3}{4}q\frac{f_{xx}}{f_x}^2 + \frac{q_x f_{xx}}{f_x} - 2r_x f_x \\ & - \frac{1}{6}\lambda q, \end{aligned} \tag{106}$$

$$r_t = \frac{3}{2}\frac{r_x f_{xxx}}{f_x} - r_{xxx} - \frac{3}{4}\frac{r_x f_{xx}^2}{f_x^2} + \frac{1}{2}\lambda, \tag{107}$$

$$\begin{aligned} s_{xx} = {} & \frac{f_{xx}s_x}{f_x} + \left(\frac{1}{2}\frac{f_{xxx}}{f_x} - \frac{3}{4}\frac{f_{xx}^2}{f_x^2} - \frac{1}{6}\lambda\right) s \\ & + 4f_x(fr_x - q_x), \end{aligned} \tag{108}$$

then

$$A'(\epsilon) = -\frac{1}{2}\frac{\epsilon^2 f_x^2}{(1+\epsilon f)^2} + \frac{1}{2}\frac{\epsilon f_{xx}}{(1+\epsilon f)} + A, \tag{109}$$

$$\begin{aligned} A'(\epsilon) = {} & -\frac{1}{2}\frac{\epsilon^2 f_x^2}{(1+\epsilon f)^2} + \frac{1}{2}\frac{\epsilon f_{xx}}{(1+\epsilon f)} + \frac{1}{2}\frac{\epsilon^2 s}{(1+\epsilon f)} \\ & + \epsilon q + A, \end{aligned} \tag{110}$$

are P_sT_d invariant solution and P_sT_d symmetry breaking solution of the AB-KdV equation respectively.

Theorem 3 can be proved true by direct substituting (109) and (110) into the AB-KdV system (2) using the Shwarzian KdV equation (104) and the anti-symmetric solutions of (105)-(108).

7. Summary and discussion

In summary, a special AB-KdV system is directly obtained from KdV equation to describe two-place physical events. The AB-KdV system possessing P_sT_d symmetry means the AB-KdV system is invariant under the transformation $\{x \to -x + x_0,\ t \to -t + t_0\}$. The AB-KdV system is nonlocal and it can be used to describe two-place physical events.

We study the exact solutions of the AB-KdV system, including P_sT_d invariant and P_sT_d symmetric breaking solutions with different methods. The P_sT_d invariant solutions, such as multiple soliton solutions, symmetry reduction solutions are obtained from the KdV equation by the P_sT_d principle. The P_sT_d symmetric breaking solutions are obtained by the help of a coupled KdV system. The solutions, such as single soliton solutions, singular soliton solutions show rich structures of the AB-KdV system. And we believe the methods can be used to other AB systems.

Also, a special Bäcklund transformation related to residual symmetry is presented via the localization of the residual symmetry to find interactive solutions between the solitons and other types of the AB-KdV system.

Acknowledgements

The authors are indebt to thank Prof. Y. Chen and B. Li for their helpful discussions. The work was supported by NNSFC (Nos. 11175092, 11435005 and 11675084), Ningbo Natural Science Foundation (No. 2015A610159) and granted by the Opening Project of Zhejiang Provincial Top Key Discipline of Physics Sciences in Ningbo University (No. xkzwl1502). And the authors were sponsored by K. C. Wong Magna Fund in Ningbo University.

[1] M. J. Ablowitzl and Z. H. Musslimani, Phys. Rev. Lett. **110**, 064105 (2013).
[2] C. Q. Song, D. M. Xiao and Z. N. Zhu, Commun. Nonlinear. Sci. Numer. Simulat. **47**, 1 (2017).
[3] M. J. Ablowitzl and Z. H. Musslimani, Nonlinearity **29**, 915 (2016).
[4] J. L. Ji and Z. N. Zhu, *Soliton solutions of an integrable nonlocal modified Korteweg-de Vries equation through inverse scattering transform*, arXiv: 1603.03994.nlin.SI (2016).
[5] M. J. Ablowitzl and Z. H. Musslimani, Phys. Rev. E **90**, 032912 (2014).
[6] M. Dimakos and A. S. Fokas, J. Math. Phys. **54**, 081504 (2014).
[7] A. S. Fokas, Phys. Rev. Lett. **96**, 190201 (2006).
[8] A. S. Fokas, Nonlinearity **29**, 319 (2016).
[9] C. M. Bender, Rep. Prog. Phys. **70**, 947 (2007).
[10] K. G. Markis, R. Ei-Ganainy, D. N. Christodoulidues, and Z. H. Musslimanni, Phys. Rev. Lett. **100**, 103904 (2008).
[11] C. M. Bender and S. P. Klevansky, Phys. Rev. Lett. **105**, 031601 (2010).

[12] Z. Lin, J. Schindler, F. M. Ellis, and T. Kottos, Phys. Rev. A **85**, 050101 (2012).
[13] T. A. Gadzhimuradov and A. M. Agalarov, Phys. Rev. A **93**, 062124 (2016).
[14] S. Y. Lou, *Alice-Bob systems, $P_s - T_d - C$ principles and multi-soliton solutions*, arXiv: 1603. 03975v2. nlin. SI, (2016).
[15] S. Y. Lou, *Alice-Bob Physics, coherent solutions of nonlocal KdV systems*, arXiv: 1606. 03154v1. nlin. SI, (2016).
[16] R. Hirota, Phys. Rev. Lett. **27**, 1192 (1971).
[17] S. Y. Lou, Stud. Appl. Math., **134**, 372 (2015).
[18] H B Lan and K L wang, J. Phys. A: Math. Gen., **23**, 4097 (1990).
[19] S. Y. Lou and G. J. Ni, J. Math. Phys., **30**,1614 (1989).
[20] S. Lie, *Vorlesungen über Differentialgleichungen mit Bekannten Infinitesimalen Transformationen*, Teuber, Leipzig (1981) (1967 reprinted by Chelsea, New York)
[21] P. J. Olver, *Applicaition of Lie Groups to Differential Equation, Graduate Texts in Mathematics*, 2nd ed., Springer-Verlag, New York (1993).
[22] G. W. Bluman and S. Kumei, *Symmetries and Differential Equation*, Appl. Math. Sci., Springer-Verlag, Berlin (1989).
[23] Gao X N, Lou S Y and Tang X Y, J. High Energy Phys. **30**, 2201 (2013).
[24] C. L. Chen, S. Y. Lou, Commun. Theor. Phys. **61**, 545 (2014).
[25] Y. Jin, M. Jia and S. Y. Lou, Commun. Theor. Phys. **58**, 795 (2012).